

University of Groningen

## Draft Genome Sequences of Five Spore-Forming Food Isolates of *Bacillus pumilus*

de Jong, Anne; van Heel, Auke J; Montalban-Lopez, Manuel; Krawczyk, Antonina O; Berendsen, Erwin M; Wells-Bennik, Marjon; Kuipers, Oscar P

*Published in:*  
Genome Announcements

*DOI:*  
[10.1128/genomeA.01539-14](https://doi.org/10.1128/genomeA.01539-14)

**IMPORTANT NOTE:** You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

*Document Version*  
Publisher's PDF, also known as Version of record

*Publication date:*  
2015

[Link to publication in University of Groningen/UMCG research database](#)

### *Citation for published version (APA):*

de Jong, A., van Heel, A. J., Montalban-Lopez, M., Krawczyk, A. O., Berendsen, E. M., Wells-Bennik, M., & Kuipers, O. P. (2015). Draft Genome Sequences of Five Spore-Forming Food Isolates of *Bacillus pumilus*. *Genome Announcements*, 3(2), 1-2. [e01539-14]. <https://doi.org/10.1128/genomeA.01539-14>

### **Copyright**

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

### **Take-down policy**

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

*Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.*

# Draft Genome Sequences of Five Spore-Forming Food Isolates of *Bacillus pumilus*

Anne de Jong,<sup>a,c</sup> Auke J. van Heel,<sup>a</sup> Manuel Montalban-Lopez,<sup>a</sup> Antonina O. Krawczyk,<sup>a,c</sup> Erwin M. Berendsen,<sup>a,b,c</sup> Marjon Wells-Bennik,<sup>b,c</sup> Oscar P. Kuipers<sup>a,c</sup>

Molecular Genetics, University of Groningen, Groningen, the Netherlands<sup>a</sup>; NIZO food research, Ede, the Netherlands<sup>b</sup>; Top Institute Food and Nutrition (TIFN), Wageningen, the Netherlands<sup>c</sup>

Here, we report the draft genome sequences of five food isolates of *Bacillus pumilus*, a spore-forming Gram-positive bacterium.

Received 15 January 2015 Accepted 2 February 2015 Published 12 March 2015

**Citation** de Jong A, van Heel AJ, Montalban-Lopez M, Krawczyk AO, Berendsen EM, Wells-Bennik M, Kuipers OP. 2015. Draft genome sequences of five spore-forming food isolates of *Bacillus pumilus*. *Genome Announc* 3(2):e01539-14. doi:10.1128/genomeA.01539-14.

**Copyright** © 2015 de Jong et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 3.0 Unported license](https://creativecommons.org/licenses/by/3.0/).

Address correspondence to Oscar P. Kuipers, o.p.kuipers@rug.nl.

*Bacillus pumilus* is a Gram-positive rod-shaped spore-forming soil bacterium. *B. pumilus* and its spores are commonly resistant to extreme environmental conditions (1, 2). Therefore, *B. pumilus* contamination in industrial settings can be persistent. *B. pumilus* is also used beneficially in the production of industrially relevant compounds, such as xylanases (3), lipases (4), and proteases (5).

The five *B. pumilus* strains used in this study were grown overnight in 10 ml of brain heart infusion (BHI) broth (Difco) at 37°C and were harvested at the exponential growth phase after reinoculation. Following centrifugation, the cell pellet was resuspended in SET buffer (75 mM NaCl, 25 mM EDTA, 20 mM Tris-HCl [pH 7.5]) and incubated with lysozyme (2 mg/ml) and RNase (0.4 mg/ml) for 30 min at 37°C. Subsequently, the sample was treated with SDS (1% final concentration) and proteinase K (0.5 mg/ml) at 55°C for 60 min. Genomic DNA was extracted from the lysate with phenol-chloroform, precipitated with isopropanol and sodium acetate (300 mM), and dissolved in Tris-EDTA (TE) buffer. The isolated DNA was sheared to 500-bp fragments in a Covaris (KBiosciences) ultrasonicator device for preparing the next-generation sequencing (NGS) library using the paired-end NEBNext Ultra DNA library prep kit for Illumina. The libraries were 101 bases paired-end sequenced on an Illumina HiSeq 2000 by multiplexing 12 samples per flow cell. Velvet (6), in combination with VelvetOptimiser (<https://github.com/Victorian-Bioinformatics-Consortium/VelvetOptimiser>), was used to perform a *de novo* paired-end assembly on each of the five genomes, resulting in the draft genome sequences (Table 1). An-

notation of the genomes was done using the following steps: (i) the scaffolds were uploaded to the RAST server (7) and automatically annotated using the SEED-based method on this server, (ii) the resulting annotated scaffolds were mapped using CONTIGuator (8) to their closest neighbor (as identified by RAST) to generate the pseudogenome, (iii) locus tags were added to each feature using an in-house-developed perl script, according to the NCBI standard, (iv) BAGEL3 (9) was used to find and annotate bacteriocin gene clusters, and (v) the protein annotation was extended using InterProScan (10)].

**Nucleotide sequence accession numbers.** The genome sequence of the five annotated strains have been deposited as whole-genome shotgun projects at DDBJ/EMBL/GenBank under the accession numbers listed in Table 1.

## ACKNOWLEDGMENTS

We thank the NGS sequence facility of the University Medical Center of Groningen (UMCG) for performing the sequencing of the strains. We also thank the Top Institute for Food and Nutrition for contributing to the funding of the project in theme 3: Safety and Preservation.

## REFERENCES

- Link L, Sawyer J, Venkateswaran K, Nicholson W. 2004. Extreme spore UV resistance of *Bacillus pumilus* isolates obtained from an ultraclean spacecraft assembly facility. *Microb Ecol* 47:159–163. <http://dx.doi.org/10.1007/s00248-003-1029-4>.
- Kempf MJ, Chen F, Kern R, Venkateswaran K. 2005. Recurrent isolation of hydrogen peroxide-resistant spores of *Bacillus pumilus* from a spacecraft assembly facility. *Astrobiology* 5:391–405. <http://dx.doi.org/10.1089/ast.2005.5.391>.

TABLE 1 Overview of the five *B. pumilus* strains in NCBI BioProject PRJNA270572

Strain	Accession no.	Isolation source
R1 B4107	JXCK00000000	Reisentopf with chicken
B4127	JXCL00000000	Cereals, cereal products, breads, pasta, and pastries
B4129	JXCM00000000	Milk, dairy products, and desserts
B4133	JXCN00000000	Cereals, cereal products, breads, pasta, and pastries
B4134	JXCO00000000	Vegetables and fermented vegetable products

3. Battan B, Sharma J, Dhiman SS, Kuhad RC. 2007. Enhanced production of cellulase-free thermostable xylanase by *Bacillus pumilus* ASH and its potential application in paper industry. *Enzyme Microb Technol* 41: 733–739. <http://dx.doi.org/10.1016/j.enzmictec.2007.06.006>.
4. Kim HK, Choi HJ, Kim MH, Sohn CB, Oh TK. 2002. Expression and characterization of Ca<sup>2+</sup>-independent lipase from *Bacillus pumilus* B26. *Biochim Biophys Acta* 1583:205–212. [http://dx.doi.org/10.1016/S1388-1981\(02\)00214-7](http://dx.doi.org/10.1016/S1388-1981(02)00214-7).
5. Huang Q, Peng Y, Li X, Wang H, Zhang Y. 2003. Purification and characterization of an extracellular alkaline serine protease with dehairing function from *Bacillus pumilus*. *Curr Microbiol* 46:169–173. <http://dx.doi.org/10.1007/s00284-002-3850-2>.
6. Zerbino DR, Birney E. 2008. Velvet: algorithms for *de novo* short read assembly using de Bruijn graphs. *Genome Res* 18:821–829. <http://dx.doi.org/10.1101/gr.074492.107>.
7. Aziz RK, Bartels D, Best AA, DeJongh M, Disz T, Edwards RA, Formsma K, Gerdes S, Glass EM, Kubal M, Meyer F, Olsen GJ, Olson R, Osterman AL, Overbeek RA, McNeil LK, Paarmann D, Paczian T, Parrello B, Pusch GD, Reich C, Stevens R, Vassieva O, Vonstein V, Wilke A, Zagnitko O. 2008. The RAST server: Rapid Annotations using Subsystems Technology. *BMC Genomics* 9:75. <http://dx.doi.org/10.1186/1471-2164-9-75>.
8. Galardini M, Biondi EG, Bazzicalupo M, Mengoni A. 2011. CONTIGuator: a bacterial genomes finishing tool for structural insights on draft genomes. *Source Code Biol Med* 6:11. <http://dx.doi.org/10.1186/1751-0473-6-11>.
9. van Heel AJ, de Jong A, Montalbán-López M, Kok J, Kuipers OP. 2013. BAGEL3: automated identification of genes encoding bacteriocins and (non-) bactericidal posttranslationally modified peptides. *Nucleic Acids Res* 41:W448–W453. <http://dx.doi.org/10.1093/nar/gkt391>.
10. Quevillon E, Silventoinen V, Pillai S, Harte N, Mulder N, Apweiler R, Lopez R. 2005. InterProScan: protein domains identifier. *Nucleic Acids Res* 33:W116–W120. <http://dx.doi.org/10.1093/nar/gki442>.